Evaluation and management of adult idiopathic intracranial hypertension

Susan P Mollan,^{1,2} Catherine Hornby,^{1,3} James Mitchell,^{1,3,4} Alexandra J Sinclair^{1,2,3,4}

¹Metabolic Neurology, Institute of Metabolism and Systems Research, University of Birmingham, Birmingham, UK

²Birmingham Neuro-Ophthalmology, Queen Elizabeth Hospital, Birmingham, UK

³Centre for Endocrinology, Diabetes and Metabolism, Birmingham Health Partners, Birmingham, UK

⁴Department of Neurology, University Hospitals Birmingham, Queen Elizabeth Hospital, Birmingham, UK

Correspondence to

Miss Susan P Mollan, Birmingham Neuro-Ophthalmology, Queen Elizabeth Hospital, Birmingham B15 2BW, UK; soozmollan@doctors.org.uk

Accepted 15 July 2018
Published Online First
28 August 2018

ABSTRACT

This paper summarises the first consensus guidelines for idiopathic intracranial hypertension as an infographic. Following a systematic literature review, a multidisciplinary specialist interest group met and established questions relating to population, interventions, controls and outcomes (PICO). A survey was sent to doctors who manage idiopathic intracranial hypertension (IIH) regularly. Statements were reviewed by national professional bodies, specifically the Association of British Neurologists, British Association for the Study of Headache, the Society of British Neurological Surgeons and the Royal College of Ophthalmologists and by international experts. Key areas are represented based on the guideline, namely: (1) investigation of papilloedema and diagnosis of IIH; (2) management strategies; and (3) investigation and management of acute exacerbation of headache in established IIH. We present an infographic as an aide-mémoire of the first consensus guidelines for IIH.

IIH is commonly associated with obesity, younger age and females.^{1 2} Patients present acutely to many different specialities and often have multiple acute visits through the course of their disease. The investigation and management of IIH is complex involving many specialities.³ This infographic summarises three key pathways based on the recommendations of a multidisciplinary, patient-involving and multiprofessional specialist interest group on the investigation and management of IIH.⁴

The basis of the specialist interest group included representation from neurology, neurosurgery, neuroradiology, ophthalmology, nursing, primary care doctors and patient representatives. Questions on PICO were defined and through a large Delphi group exercise; expertise was captured from a wide-reaching group of

clinicians, thus reflecting practice from across the UK and internationally. The statements were then critically reviewed by key opinion leaders and by Association of British Neurologists, British Association for the Study of Headache, the Society of British Neurological Surgeons and the Royal College of Ophthalmologists. This is the first consensus guidance for optimal management of IIH.⁴

Identification of papilloedema can be challenging, and clinicians should be aware of the differential diagnosis of pseudopapilloedema (figure 1). Once papilloedema is confirmed, it requires urgent investigations, including lumbar puncture, where the patient experience could be greatly improved.⁵ Symptoms of IIH are not pathognomonic, and hence it is essential to apply the diagnostic criteria, including excluding secondary causes, for a definite diagnosis.⁴ The lumbar puncture opening pressure was one key area of debate. Within the wider Delphi group, it was clear that there is a 'grey zone' of lumbar puncture opening pressures between 25 cm cerebrospinal fluid (cmCSF) and 30 cmCSF, as to what each expert considered to be pathological, and this is reflected within the infographic thermometer for lumbar puncture opening pressure (figure 1).

Principles of management need to address both the rapidity of the disease that may lead to visual loss in some and require surgical intervention and the morbidity of the headache that can develop in the majority, which substantially affects the quality of life.⁶ Weight loss is currently the only established disease-modifying therapy⁷ and is notoriously difficult to achieve and maintain.

Evaluation of the headache phenotype is essential to target treatment and to help identify medication-overuse headache. Where there are features of migraine, topiramate may be the first line in treatment,

© Author(s) (or their employer(s)) 2018. Re-use permitted under CC BY. Published by BMJ.

To cite: Mollan SP, Hornby C, Mitchell J, et al. *Pract Neurol* 2018;**18**:485–488.

Investigation of Papilloedema

Papilloedema identified

Assess vision

Record:
Visual acuity
Pupil examination
Formal visual fields
Dilated fundoscopy
*For experienced clinician review - high risk of misdiagnosis

Check BP
Exclude malignant hypertension
>180/120mmHg²

URGENT Brain imaging within 24 hours (CT/MRI) + Venography essential

IIH imaging features:

- Partially/empty sella
- Increased tortuosity of optic nerve
- Enlarged optic nerve sheath
- Flattened posterior globe/sclera
- Intracanal protrusion of optic nerve head
- Attenuation of cerebrosinus sinuses e.g. transverse sinus stenosis

Other causes of bilateral disc swelling

- Infective
- Inflammatory
- Hypertensive crisis
- Toxic
- Metabolic
- Hereditary
- Compressive (Uncommon vascular)

Pseudopapilloedema

- Congenitally anomalous discs
- Optic nerve head drusen
- Hypermetropic discs
- Myopic discs
- Myelinated nerve fibres

Exclude secondary causes

- Anaemia
- Venous thrombosis
- Drugs
- Endocrine disease/syndromes
- Drugs eg fluoroquinolones, tetracyclines, vitamin A analogues, withdrawal of long term glucocorticoids

✓ No lesion identified

Lumbar puncture

CSF opening pressure of >25cmCSF; normal constituents

cmCSF

Increasing likelihood of being pathological

Pressure in 'grey zone' may be normal for some individuals

Diagnostic criteria cut off

Idiopathic Intracranial Hypertension

Symptoms

Frequency of symptoms reported	Headache	Neck pain
>65% ★★★	★★★	★★★
40-65% ★★	★★★	★★★
<30% ★★	★★★	★★★

Visual obscurations ★★★

Pulsatile tinnitus ★★

Back pain ★★

Dizziness ★★

Neck pain ★★★

Blurred vision ★★★

Cognitive disturbance ★★

Radicular pain ★★

Diplopia ★★

Diagnostic criteria

- Papilloedema
- Normal neurological examination (except sixth nerve palsy)
- Neuroimaging: normal brain parenchyma. Venous thrombosis excluded
- Normal CSF constituents
- Elevated LP pressure >25cmCSF

Management Strategies

Principles of management

- Protect vision**
- Manage underlying disease**
- Reduce headache morbidity**

Atypical IIH
Patients who are not female, or not of child bearing age or who have a BMI below 30kg/m²

Typical IIH
Patients who are female, of child bearing age and who have BMI >30kg/m²

Fulminant IIH
Patients with declining visual function within 4 weeks of diagnosis of IIH

Consider medical therapy with acetazolamide

Temporising lumbar drain if surgery planned >24 hours

CSF diversion OR Optic nerve sheath fenestration

Weight management

Headache assessment + Ongoing visual assessment

Significant deterioration of visual function

LP

If pathologically high

If not re-evaluate

VISION THREATENED

Acute Exacerbation of Headache in IIH

Emergency Room Attendance due to headache

Mandatory assessment of papilloedema

Papilloedema

No papilloedema

Assess vision
As above

Exclude secondary causes and red flags e.g. meningitis and bleed

Patient involvement

IIH headache flare-up

- × No papilloedema
- × No red flags or other secondary causes
- × No imaging required
- × No LP required

Address medication overuse

- × Simple analgesics >15 days/month
- × Opiates/triptans >10 days/month

Evaluate headache phenotype

- Elevated ICP
- Migraine
- Medication overuse
- CSF shunt in situ
- ? Low pressure
- ? High pressure
- ? Normal

Treat migraine headache

Acute migraine attack

- ✓ Triptan
- ✓ NSAID/paracetamol
- ✓ Anti-emetic
- × No opiates

Maximum 10 times per month

Migraine prevention

- ✓ 3 month trial
- ✓ Titrate up slowly
- ✓ Reach a therapeutic dose

Options

- Topiramate
- Candesartan
- Botox (as per NICE)
- +/- tricyclics
- +/- propranolol
- +/- flunarizine

Caution: ↑ weight, ↓ mood

1. Mollan SP, Davis B, Silver NC, Shaw S, Maluco C, Wakerley BP, Kfirian A, Chavda S, Ramalingam S, Edwards J, Hemmings K, Williamson M, Burton MA, Hassan-Smith G, Digre K, Liu GT, Jensen RH, Sinclair AJ. Idiopathic intracranial hypertension: International consensus guidelines. *J Neurol Neurosurg Psychiatry*. 2019;90(10):1109-1117. doi:10.1136/npp.2019.227000.

2. Whetton PC, Carey RM, Aronow WS, et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APHA/ASH/ASPC/NMA/PCNA guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension*. 2017;70(5):e133-e115. doi:10.1161/HYPERTENSION.2017.04.016.

and recent evidence indicates that it has a significant intracranial pressure-lowering effect in rodents.⁸ Acute exacerbation of headache often leads to reinvestigation with lumbar puncture, and the collective expert opinion reflected that lumbar puncture provides only temporary relief, can lead in some to longer term complications⁹ and exacerbation of headache.¹⁰ In those with acute exacerbation of headache, optic nerve examination is essential, and in those found not to have papilloedema, investigation with lumbar puncture and brain imaging is not required, so long as no other secondary causes of headache are suspected. The infographic illustrates the management of acute exacerbation of headache in IIH (figure 1).

Horizon scanning for IIH shows that research is active and that metabolic concepts may potentially provide more understanding of the cause and provide evidence for innovative therapeutic opportunities.¹¹ A phase 2 randomised control trial with the first novel drug treatment for IIH has finished recruitment¹²; a phase 3 randomised control trial investigating the best method for weight loss is underway¹³; other surgical trials are in planning.

This infographic highlights three areas that are covered by the consensus guideline for adult IIH, which are: (1) investigation of papilloedema and diagnosis of IIH; (2) management strategies; and (3) investigation and management of acute exacerbation of headache in established IIH⁴ (figure 1).

Key points

- ▶ Cerebral venography is an essential part of the work-up to exclude venous sinus thrombosis as a cause of papilloedema.
- ▶ Lumbar puncture opening pressure forms part of the diagnostic criteria; however, most clinicians feel there is a 'grey zone' between 25 cmCSF and 30 cmCSF, which may not be pathological.
- ▶ Those with fulminant or precipitous visual decline need urgent surgical treatment, preferably with a ventriculoperitoneal shunt.
- ▶ All patients diagnosed with idiopathic intracranial hypertension need sensitive and appropriate discussion regarding weight loss (the only disease-modifying treatment).
- ▶ Those with acute exacerbation of headache do not require further neuroimaging or repeat lumbar puncture, unless there are red flag symptoms/signs of infection, or papilloedema with precipitous visual decline.

Correction notice This article has been corrected since it was published Online First. The provenance and peer review disclaimer incorrectly said the paper was not commissioned.

Acknowledgements We acknowledge all those who contributed to the surveys and critiqued the document in the wider Delphi group. The specialist interest group members and international

key opinion leaders: Brendan Davis (University Hospital North Midlands NHS Trust), Nick Silver (The Walton Centre NHS Foundation Trust), Simon Shaw S (University Hospital North Midlands NHS Trust), Conor Mallucci (The Walton Centre NHS Foundation Trust and Alder Hey Children's NHS Foundation Trust), Ben Wakerley (Gloucestershire Hospitals NHS Foundation Trust), Anita Krishnan (The Walton Centre NHS Foundation Trust), Swarup Chavda (University Hospitals Birmingham NHS Foundation Trust), Sateesh Ramalingam (University Hospitals Birmingham NHS Foundation Trust), Julie Edwards (University Hospitals Birmingham NHS Foundation Trust and Sandwell and West Birmingham NHS Trust), Krystal Hemmings (IIH UK Charity), Shelly Williamson (IIH UK Charity), Mike Burdon (University Hospitals Birmingham NHS Foundation Trust), Ghaniah Hassan-Smith G (University of Birmingham), Kathleen Digre (Moran Eye Center, University of Utah) Grant Liu (Children's Hospital of Philadelphia and Hospital of the University of Pennsylvania), Rigmor Jensen (Danish Headache Centre, Copenhagen). Collaborators: Jane Anderson, (Cambridge University Hospitals NHS Foundation Trust), Peter Goadsby (King's College Hospital), Tim Matthews (University Hospitals Birmingham NHS Foundation Trust) and Jan Hoffman (University Medical Centre Hamburg).

Collaborators Brendan Davis (University Hospital North Midlands NHS Trust), Nick Silver (The Walton Centre NHS Foundation Trust), Simon Shaw (University Hospital North Midlands NHS Trust), Conor Mallucci (The Walton Centre NHS Foundation Trust and Alder Hey Children's NHS Foundation Trust), Ben Wakerley (Gloucestershire Hospitals NHS Foundation Trust), Anita Krishnan (The Walton Centre NHS Foundation Trust), Swarup Chavda (University Hospitals Birmingham NHS Foundation Trust), Sateesh Ramalingam (University Hospitals Birmingham NHS Foundation Trust), Julie Edwards (University Hospitals Birmingham NHS Foundation Trust and Sandwell and West Birmingham NHS Trust), Krystal Hemmings (IIH UK Charity), Shelly Williamson (IIH UK Charity), Mike Burdon (University Hospitals Birmingham NHS Foundation Trust), Ghaniah Hassan-Smith (University of Birmingham), Kathleen Digre (Moran Eye Center, University of Utah) Grant Liu (Children's Hospital of Philadelphia and Hospital of the University of Pennsylvania), Rigmor Jensen (Danish Headache Centre, Copenhagen), Jane Anderson, (Cambridge University Hospitals NHS Foundation Trust), Peter Goadsby (King's College Hospital), Tim Matthews (University Hospitals Birmingham NHS Foundation Trust) and Jan Hoffman (University Medical Centre Hamburg).

Contributors SPM and AJS drafted the infogram. CH executed the drawing of the infogram. SPM wrote the initial draft. JM and AJS critically reviewed the paper and infogram.

Funding AJS is funded by an NIHR Clinician Scientist Fellowship (NIHR-CS-011-028) and by the Medical Research Council, UK (MR/K015184/1).

Competing interests None declared.

Patient consent Not required.

Provenance and peer review Commissioned; externally peer reviewed by Joseph Anderson, Cardiff, UK and Mark Lawden, Leicester, UK. The guideline was also reviewed, critiqued and supported by the following professional bodies: The Association of British Neurologists, the Society of British Neurological Surgeons, the Royal College of Ophthalmologists and the British Association for the Study of Headache.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any

purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

REFERENCES

- 1 Mollan SP, Ali F, Hassan-Smith G, *et al*. Evolving evidence in adult idiopathic intracranial hypertension: pathophysiology and management. *J Neurol Neurosurg Psychiatry* 2016;87:982–92.
- 2 Markey KA, Mollan SP, Jensen RH, *et al*. Understanding idiopathic intracranial hypertension: mechanisms, management, and future directions. *Lancet Neurol* 2016;15:78–91.
- 3 Mollan SP, Markey KA, Benzimra JD, *et al*. A practical approach to, diagnosis, assessment and management of idiopathic intracranial hypertension. *Pract Neurol* 2014;14:380–90.
- 4 Mollan SP, Davies B, Silver NC, *et al*. Idiopathic intracranial hypertension: consensus guidelines on management. *J Neurol Neurosurg Psychiatry* 2018.
- 5 Scotton WJ, Mollan SP, Walters T, *et al*. Characterising the patient experience of diagnostic lumbar puncture in idiopathic intracranial hypertension: a cross-sectional online survey. *BMJ Open* 2018;8:e020445.
- 6 Mulla Y, Markey KA, Woolley RL, *et al*. Headache determines quality of life in idiopathic intracranial hypertension. *J Headache Pain* 2015;16:45.
- 7 Sinclair AJ, Burdon MA, Nightingale PG, *et al*. Low energy diet and intracranial pressure in women with idiopathic intracranial hypertension: prospective cohort study. *BMJ* 2010;341:c2701.
- 8 Scotton WJ, Botfield HF, Westgate CS, *et al*. Topiramate is more effective than acetazolamide at lowering intracranial pressure. *Cephalalgia* 2018;1:033310241877645.
- 9 Engelborghs S, Niemantsverdriet E, Struyfs H, *et al*. Consensus guidelines for lumbar puncture in patients with neurological diseases. *Alzheimers Dement* 2017;8:111–26.
- 10 Yiangou A, Mitchell J, Markey KA, *et al*. Therapeutic lumbar puncture for headache in idiopathic intracranial hypertension: Minimal gain, is it worth the pain? *Cephalalgia* 2018;1:033310241878219.
- 11 Hornby C, Mollan SP, Botfield H, *et al*. Metabolic concepts in idiopathic intracranial hypertension and their potential for therapeutic intervention. *J Neuroophthalmol* 2018. doi: 10.1097/WNO.0000000000000684. [Epub ahead of print 6 Jul 2018].
- 12 Markey KA, Ottridge R, Mitchell JL, *et al*. Assessing the Efficacy and Safety of an 11 β -Hydroxysteroid Dehydrogenase Type 1 Inhibitor (AZD4017) in the Idiopathic Intracranial Hypertension Drug Trial, IIH:DT: clinical methods and design for a phase II randomized controlled trial. *JMIR Res Protoc* 2017;6:e181.
- 13 Ottridge R, Mollan SP, Botfield H, *et al*. Randomised controlled trial of bariatric surgery versus a community weight loss programme for the sustained treatment of idiopathic intracranial hypertension: the Idiopathic Intracranial Hypertension Weight Trial (IIH:WT) protocol. *BMJ Open* 2017;7:e017426.